



C. INTRAOCULAR PRESSURE			
O.D.	O.S.	TEST METHOD USED	TIME SINCE LAST MEDICATION.
NOTE - Pressures should NOT be taken within 2 hours after use of medication unless 10.B. is completed.			
11. FIELD OF VISION - Record physiological and any pathological peripheral or central visual field losses from a perimeter and/or tangent screen using white test object. - <i>FORWARD CHARTS.</i>			
A. DID EXAMINE WEAR GLASSES OR CONTACT LENSES DURING TEST? (Specify which)		B. SIZE OF TEST OBJECT USED WITH TANGENT SCREEN	
12 VISUAL ACUITY - Record ( <i>Use Snellen linear values</i> )			
A. DISTANT	TEST METHOD USED	UNCORRECTED O.D.    O.S.    O.U.	CORRECTED O.D.    O.S.    O.U.
B. INTERMEDIATE (32 inches)	TEST METHOD USED	UNCORRECTED O.D.    O.S.    O.U.	CORRECTED O.D.    O.S.    O.U.
C. NEAR	TEST METHOD USED	UNCORRECTED O.D.    O.S.    O.U.	CORRECTED O.D.    O.S.    O.U.
D. IMPORTANT - If correction is needed and there is inability to correct either eye to 20/20 or better, give reasons.			
13. PRESENT CONDITION			
A. DOES AIRMAN WEAR		O.D.	O.S.
		SPHERE-CYLINDER-AXIS	SPHERE-CYLINDER-AXIS
GLASSES	CONTACT LENSES		
14. PUPILS - Statement of relative size and reaction of the pupils to accommodation and light, with special reference to any disease process, healed or active.			
15. OPHTHALMOSCOPIC - Describe any variation from normal in either eye on fundoscopic examinations, with special reference to any disease process, healed or active.			
16. SLIT LAMP - Record results of slit lamp examination of each eye where indicated.			
17. FUSION - Estimate fusion ability and state methods used in examination.			
18A. TYPED NAME AND ADDRESS OF OPHTHALMOLOGIST		18B. SIGNATURE OF OPHTHALMOLOGIST	